

Adefin XL

contains the active ingredient nifedipine

CONSUMER MEDICINE INFORMATION

What is in this leaflet

This leaflet answers some common questions about ADEFIN XL tablets.

It is not to be used in relation to any other product, which may also contain the same active ingredient.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking ADEFIN XL against the benefits expected for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Please read this leaflet carefully and keep it with your medicine.

You may need to read it again.

What ADEFIN XL is used for

ADEFIN XL is used to:

- * treat high blood pressure
- * prevent chronic stable angina, a type of angina.

ADEFIN XL is not used to relieve sudden attacks of angina.

ADEFIN XL belongs to a group of medicines called calcium channel blockers.

Calcium channel blockers work by opening up blood vessels in the body to lower blood pressure and improve the supply of blood and oxygen to the heart.

ADEFIN XL tablets are specially designed with a strong outer shell. This outer shell allows the medicine to be released slowly into the body over a longer period of time after it is taken. The outer shell is excreted or passed out in the faeces (or bowel motions), as it does not dissolve in the body. Therefore, it is normal to find a "tablet-like" object in the faeces.

Ask your doctor if you have any questions about why ADEFIN XL has been prescribed for you.

Your doctor may have prescribed it for another reason.

This medicine is only available with a doctor's prescription.

Before you take ADEFIN XL

When you must not take it

Do not take ADEFIN XL if you are allergic to:

- * medicines containing nifedipine (eg. Adalat OROS, Adefin)
- * medicines similar to nifedipine such as amlodipine (Norvasc), felodipine (Plendil, Felodur), lercanidipine (Zanidip), nimodipine (Nimotop)
- * any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include skin rash, itching or hives; swelling of the face, lips or tongue which may cause

difficulty in swallowing or breathing; wheezing or shortness of breath.

Do not take ADEFIN XL if you are pregnant or plan to become pregnant.

It may affect your developing baby if you take it during pregnancy.

Do not take ADEFIN XL if you are breastfeeding or plan to breastfeed.

ADEFIN XL passes into breast milk and may affect your baby.

Do not take ADEFIN XL if you have had a heart attack in the last 8 days.

Do not take ADEFIN XL if you are in cardiogenic shock (very low blood pressure due to a failing heart).

Symptoms of cardiogenic shock include rapid, shallow breathing, cold, clammy skin, a rapid, weak pulse, dizziness, weakness and fainting.

Do not take ADEFIN XL if you have a Kock Pouch or ileostomy.

Do not take ADEFIN XL if you are taking any medicine containing rifampicin (Rifadin, Rimycin).

Do not take ADEFIN XL after the expiry date (EXP) printed on the pack has passed.

If you take this medicine after the expiry date, it may not work as well.

Do not take ADEFIN XL if the packaging shows signs of tampering or the tablets are broken, discoloured or show any other sign of deterioration.

When to take it

The tablets should generally be taken in the morning.

ADEFIN XL can be taken either with or without a meal.

How long to take it

Keep taking ADEFIN XL for as long as your doctor recommends.

This medicine helps to control your condition, but does not cure it. It is important to keep taking your medicine even if you feel well.

If you forgot to take it

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as soon as you remember, and then go back to taking your medicine as you would normally.

Do not take a double dose to make up for the dose you missed.

This may increase the chance of you getting an unwanted side effect.

If you have missed several doses, consult your doctor.

If you are not sure what to do, check with your doctor or pharmacist.

If you have trouble remembering to take your ADEFIN XL, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or Poisons Information Centre (telephone 13 11 26), or go to Accident and Emergency at your nearest hospital, if you think that you or anyone else may have taken too much ADEFIN XL.

Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

Signs of an overdose include dizziness and fainting due to a drop in blood pressure; an irregular heart beat; shortness of breath and even loss of consciousness.

While you are taking ADEFIN XL

Things you must do

Take ADEFIN XL exactly as prescribed by your doctor.

If you become pregnant while taking ADEFIN XL, tell your doctor immediately.

It may affect your developing baby if you take it during pregnancy.

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking ADEFIN XL

Tell your doctor if you continue to have angina attacks or if they become more frequent while you are taking ADEFIN XL.

If you are about to have any blood tests or x-rays, tell your doctor that you are taking ADEFIN XL.

It may interfere with the results of some tests.

Tell all doctors, dentists and pharmacists who are treating you that you are taking ADEFIN XL.

Things you must not do

Do not take ADEFIN XL to treat any other complaints unless your doctor tells you to.

Do not give this medicine to anyone else, even if they have the same symptoms or condition as you.

Do not stop taking your medicine or lower the dosage without checking with your doctor.

Things to be careful of

Make sure you know how ADEFIN XL affects you before driving a car, operating machinery, or do anything else that could be dangerous.

ADEFIN XL may cause dizziness or fainting in some people, especially when they first start taking the medicine, change dose, or drink alcohol.

If you have angina, be careful not to overdo physical activities when you first start taking ADEFIN XL.

You may feel better when you start taking it, but you will need time to improve your physical fitness.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking ADEFIN XL.

ADEFIN XL helps most people, but it may have unwanted effects in some people.

All medicines have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of these side effects.

Do not be alarmed by the following lists of side effects.

You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- * headache, dizziness
- * constipation
- * feeling sick (nausea)
- * flushing
- * loss of energy or unusual tiredness or weakness
- * numbness in the hands and feet.

The above list includes the more common and mild side effects.

Tell your doctor as soon as possible if you notice any of the following:

- * blurred vision
- * bleeding, tender or swollen gums
- * swelling of ankles, feet or lower leg
- * painful, swollen joints or muscles
- * trembling
- * rash or blistering of the skin when exposed to sunlight
- * unusual bleeding or bruising under the skin
- * passing more urine than normal
- * breast enlargement (in men)
- * signs of frequent infections such as fever, severe chills, sore throat or mouth ulcers.

The side effects listed above could be serious. You may need medical attention.

If any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:

- * fainting
- * fast or irregular heartbeat
- * chest pain
- * signs of an allergic reaction such as rash, itching or hives on the skin; swelling of the face, lips, tongue, or other parts of the body; shortness of breath, wheezing, or trouble breathing.
- * signs of liver problems such as yellowing of the skin and/or eyes (jaundice)

These are very serious side effects. If you have these side effects you may need urgent medical attention or hospitalisation.

In a small number of cases of in vitro fertilization, medicines like nifedipine may have interfered with the normal function of sperm. This effect went away after the medicine was stopped. In those men who are taking Adefin XL and are repeatedly unsuccessful in fathering a child by in vitro fertilization, the medicine should be considered as one of the possible causes if no other explanation can be found.

Tell your doctor if you notice anything else that is making you feel unwell.

Other side effects not listed above may also occur in some people.

After taking ADEFIN XL

Tell your doctor if you experience any symptoms of bowel blockage including:

- * abdominal swelling
- * severe abdominal cramps or pain
- * vomiting, constipation.

Very rarely, the undissolved tablet shell may not pass out in the faeces and collect in the stomach. Surgical intervention may be necessary to remove these shells.

Storage

Keep ADEFIN XL where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Keep your tablets in a cool dry place, away from sunlight where the temperature stays below 25 degrees C.

Do not store ADEFIN XL or any other medicine in the bathroom or near a sink.

Do not leave ADEFIN XL in the car or on window sills.

Keep the tablets in the pack until it is time to take them.

Heat and moisture can affect ADEFIN XL tablets. Contact with moisture can release the active ingredient (nifedipine) from the tablets. To prevent this, the tablets must be kept in the original pack until immediately before use.

Disposal

If your doctor tells you to stop taking ADEFIN XL, or your tablets have passed their expiry date, ask your pharmacist what to do with any that are left over.

Product description

What it looks like

ADEFIN XL is available in two strengths:

- * ADEFIN XL 30 - round, pink coloured tablet, with "30" printed in black on one side and blank on the other side.
- * ADEFIN XL 60 - round, pink coloured tablet, with "60" printed in black on one side and blank on the other side.

Each pack contains 30 tablets.

Ingredients

The active ingredient in ADEFIN XL is nifedipine.

- * Each ADEFIN XL 30 tablet contains 30 mg of nifedipine.
- * Each ADEFIN XL 60 tablet contains 60 mg nifedipine.

ADEFIN XL also contains the following inactive ingredients:

- * polyethylene oxide
- * magnesium stearate
- * hypromellose
- * iron oxide red CI77491 (E172)
- * hydroxypropylcellulose
- * cellulose acetate
- * macrogol 3350
- * titanium dioxide (171)
- * propylene glycol.

Supplier

ADEFIN XL is supplied by:

Alphapharm Pty Limited

(ABN 93 002 359 739)

Chase Building 2

Wentworth Park Road

Glebe NSW 2037

Phone: (02) 9298 3999

www.alphapharm.com.au

Medical Information

Phone: 1800 028 365

Australian registration numbers:

ADEFIN XL 30 - AUST R 91779

ADEFIN XL 60 - AUST R 91778

This leaflet was prepared on
17 October 2005.