


APO-DIAZEPAM TABLETS**NAME OF THE MEDICINE**

Diazepam

Chemical Name: 7-chloro-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one.

Structural Formula:

Molecular Formula: $C_{16}H_{13}ClN_2O$.

Molecular weight: 284.74.

CAS registry number: 439-14-5.

DESCRIPTION

Diazepam is a benzodiazepine derivative. It is a colourless crystalline compound, insoluble in water.

PHARMACOLOGY**Pharmacokinetics****Absorption and bioavailability**

Diazepam is rapidly and completely absorbed from the gastrointestinal tract, peak plasma concentrations appearing 30-90 minutes after oral intake. The speed of onset after intramuscular administration is variable, depending on the muscle mass used and other factors.

Distribution

Diazepam is 98% protein-bound in the plasma, and is excreted mainly (about 70%) in the urine in free form or (predominantly) as conjugated metabolites. Diazepam and its metabolites cross the blood-brain and placental barriers and are also found in breast-milk.

Metabolism

It is metabolised to hydroxy-diazepam (temazepam) and nordiazepam ($t_{1/2}$ approximately 96 hours) and ultimately to oxazepam.

Elimination

The plasma concentration time curve of diazepam is biphasic, an initial rapid and extensive distribution phase with a half-life of up to 3 hours, followed by a prolonged terminal elimination phase (half-life 20 to 48 hours). The elimination half-life is 90 hours at age 80 and increased two to three fold in patients with cirrhosis (see also **PRECAUTIONS**). The elimination half-life may be prolonged in the newborn, the elderly and patients with hepatic or renal disease and it should be noted that the plasma concentration may take correspondingly longer to reach steady state.

Pharmacodynamics

Diazepam is a member of the group of classical benzodiazepines and exhibits anxiolytic, sedative, muscle relaxant and anticonvulsant effects. This is presumed to be the result of facilitating the action in the brain of gamma-aminobutyric acid, a naturally occurring inhibitory transmitter.

INDICATIONS

- The management of anxiety disorders or for the short term relief of the symptoms of anxiety. Anxiety or tension associated with the stress of everyday life usually does not require treatment with an anxiolytic.
- In acute alcohol withdrawal, diazepam may be useful in the symptomatic relief of acute agitation, tremor, impending or acute delirium tremens and hallucinosis.
- Diazepam is a useful adjunct for the relief of reflex muscle spasm due to local trauma (injury, inflammation) to muscles, bones and joints. It can also be used to combat spasticity due to upper motor neuron lesions such as cerebral palsy and paraplegia, as well as in athetosis and stiff-man syndrome.

CONTRAINDICATIONS

Diazepam is contra-indicated in patients with:

- Known hypersensitivity to benzodiazepines or any excipients in the product
- Chronic obstructive airways disease with incipient respiratory failure
- Severe respiratory insufficiency
- Severe hepatic insufficiency
- Sleep apnoea syndrome
- Myasthenia gravis
- Dependence on CNS depressants including alcohol. An exception to the latter is the management of acute withdrawal reactions.

Benzodiazepines are not recommended for the primary treatment of psychotic illness.

Benzodiazepines should not be used alone to treat depression or anxiety associated with depression as suicide may occur in such patients.

PRECAUTIONS

Patients should be advised that their tolerance for alcohol and other CNS depressants will be diminished and that these medications should either be eliminated or given in reduced dosage in the presence of diazepam.

Following the prolonged use of diazepam at therapeutic doses, withdrawal from the medication should be gradual. An individualised withdrawal timetable needs to be planned for each patient in whom dependence is known or suspected. Periods from four weeks to four months have been suggested. As with other benzodiazepines, when treatment is suddenly withdrawn, a temporary increase in sleep disturbance can occur after use of diazepam (see **PRECAUTIONS – Dependence**).

In general, benzodiazepines should be prescribed for short periods only (e.g. 2-4 weeks). Continuous long-term use of diazepam is not recommended. There is evidence that tolerance develops to the sedative effects of benzodiazepines. After as little as one week of therapy, withdrawal symptoms can appear following the cessation of recommended doses (e.g. rebound insomnia following cessation of a hypnotic benzodiazepine).

Since the tablets contain lactose, patients with rare hereditary problems of galactose intolerance (the Lapp lactase deficiency or glucose-galactose malabsorption) should not take this medicine.

Effect on Ability to Drive or Operate Machinery

Sedation, amnesia, impaired concentration and impaired muscle function may adversely affect the ability to drive or operate machinery. As with all patients taking CNS-depressant medications, patients receiving diazepam should be warned not to operate dangerous machinery or motor vehicles until it is known that they do not become drowsy or dizzy from diazepam therapy. Abilities may be impaired on the day following use.

Effect on Blood Pressure

Although hypotension has occurred rarely, diazepam should be administered with caution to patients in whom a drop in blood pressure might lead to cardiac or cerebral complications. This is particularly important in elderly patients.

Memory Impairment

Transient amnesia or memory impairment has been reported in association with the use of benzodiazepines. Anterograde amnesia may occur using therapeutic dosages, the risk increasing at higher dosages. Amnestic effects may be associated with inappropriate behaviour.

Glaucoma (Narrow Angle)

Caution should be used in the treatment of patients with acute narrow-angle glaucoma (because of atropine-like side effects).

Impaired Renal / Liver Function and Blood Dyscrasias

Patients with impaired renal or hepatic function should use benzodiazepine medication with caution and dosage reduction may be advisable. In rare instances, some patients taking benzodiazepines have developed blood dyscrasias, and some have had elevation of liver enzymes. As with other benzodiazepines, periodic blood counts and liver function tests are recommended.

Depression, Psychosis and Schizophrenia

Diazepam is not recommended as primary therapy in patients with depression and/or psychosis. In such conditions, psychiatric assessment and supervision are necessary if benzodiazepines are indicated. Benzodiazepines may increase depression in some patients and may contribute to deterioration in severely disturbed schizophrenics with confusion and withdrawal. Suicidal tendencies may be present or uncovered and protective measures may be required.

Paradoxical Reactions

Paradoxical Reactions such as restlessness, agitation, irritability, aggressiveness, delusion, nightmares, hallucinations, psychoses, inappropriate behaviour and other adverse behavioural effects, acute rage, stimulation or excitement may occur. Should such reactions occur, diazepam should be discontinued. They are more likely to occur in children and in the elderly.

Impaired Respiratory Function

Caution in the use of diazepam is recommended in patients with respiratory depression. In patients with chronic obstructive pulmonary disease, benzodiazepines can cause increased arterial carbon dioxide tension and decreased oxygen tension. A lower dose is recommended for patients with chronic respiratory insufficiency, due to the risk of respiratory depression.

Epilepsy

When diazepam is administered to persons with convulsive disorders, an increase in the frequency and/or severity of grand mal seizures may occur, necessitating increased anticonvulsant medication. Abrupt withdrawal of benzodiazepines in persons with convulsive disorders may be associated with a temporary increase in the frequency and/or severity of seizures.

Abuse

Extreme caution must be exercised in administering diazepam to individuals with a history of alcohol or drug abuse or those known to be addiction prone or those whose history suggests they may increase the dosage on their own initiative. It is desirable to limit repeat prescription without adequate medical supervision.

Dependence

The use of benzodiazepines may lead to dependence, as defined by the presence of a withdrawal syndrome on discontinuation of the drug. The risk of dependence increases with dose and duration of treatment. It is more pronounced in patients on long-term therapy and/or high dosage and particularly so in predisposed patients with a history of alcohol or drug abuse. Tolerance, as defined by a need to increase the dose in order to achieve the same therapeutic effect, seldom occurs in patients receiving recommended doses under medical supervision. Tolerance to sedation may occur with benzodiazepines, especially in those with drug seeking behaviour.

Withdrawal symptoms similar in character to those noted with barbiturates and alcohol have occurred once physical dependence to benzodiazepines has developed or following abrupt discontinuation of benzodiazepines. These symptoms range from insomnia, anxiety, dysphoria, palpitations, panic attacks, vertigo, myoclonus, akinesia, hypersensitivity to light, sound and touch, abnormal body sensations (e.g. feeling of motion, metallic taste), depersonalisation, derealisation, delusional beliefs, hyperreflexia and loss of short term memory, to a major syndrome which may include convulsions, tremor, abdominal and muscle cramps, confusional state, delirium, hallucinations, hyperthermia, psychosis, vomiting and sweating. Such manifestations of withdrawal, especially the more serious ones, are more common in patients who have received excessive doses over a prolonged period.

However, withdrawal symptoms have been reported following abrupt discontinuation of benzodiazepines taken continuously at therapeutic levels. Accordingly, diazepam should be terminated by tapering the dose to minimise occurrence of withdrawal symptoms. Patients should be advised to consult with their physician before either increasing the dose or abruptly discontinuing the medication.

Rebound phenomena have been described in the context of benzodiazepine use. Rebound insomnia and anxiety mean an increase in the severity of these symptoms beyond pre-treatment levels following cessation of benzodiazepines. Rebound phenomena in general possibly reflect re-emergence of pre-existing symptoms combined with withdrawal symptoms described earlier. Some patients prescribed benzodiazepines with very short half-lives (in the order of 2 to 4 hours) may experience relatively mild rebound symptoms in between their regular doses. Withdrawal/rebound symptoms may follow high doses for relatively short periods.

Effects on Fertility

Reproductive studies in rats showed decreases in the number of pregnancies and in the number of surviving offspring following administration of oral doses of 100 mg/kg/day (22-fold the MRHD on a body surface area basis) to both males and females prior to and during mating and throughout gestation and lactation.

Use in Pregnancy (Category C)

The safety of diazepam for use in human pregnancy has not been established. Diazepam and its metabolites readily cross the placenta. An increased risk of congenital malformation associated with the use of benzodiazepines during the first trimester of pregnancy has been suggested. Benzodiazepines should be avoided during pregnancy unless there is no safer alternative. Benzodiazepines cross the placenta and may cause hypotension, hypotonia, reduced respiratory function and hypothermia in the newborn infant. Continuous treatment during pregnancy and administration of high doses in connection with delivery should be avoided.

Withdrawal symptoms in newborn infants have been reported with this class of drugs. Special care must be taken when diazepam is used during labour and delivery, as single high doses may produce irregularities in the foetal heart rate and hypotonia, poor sucking, hypothermia and moderate respiratory depression in the neonate (floppy infant syndrome). With newborn infants it must be remembered that the enzyme system involved in the breakdown of the drug is not yet fully developed (especially in premature infants).

Category C: Drugs which, owing to their pharmacological effects, have caused or may be suspected of causing, harmful effects on the human foetus or neonate without causing malformations. These effects may be reversible. Accompanying texts should be consulted for further details.

Use in Lactation

Diazepam is excreted in human breast milk and may cause drowsiness and feeding difficulties in the infant. Breast-feeding is therefore not recommended in patients receiving diazepam.

Use in Children

Efficacy and safety of diazepam has not been established in the neonate (30 days or less in age). Prolonged central nervous system depression has been observed in neonates due to inability to transform the drug. In view of lack of adequate clinical experience chronic oral use is not recommended in children younger than 6 months.

Use in Elderly or Debilitated Patients

An increased risk of falls and fractures has been recorded in elderly benzodiazepine users.

Elderly or debilitated patients may be particularly susceptible to the sedative effects of benzodiazepines and associated giddiness, ataxia and confusion, which may increase the risk of a fall.

Concomitant use of barbiturates, alcohol, or other central nervous system depressants increases depression with increased risk of apnoea.

Lower doses should be used for elderly and debilitated patients.

Carcinogenicity

The carcinogenic potential of oral diazepam has been studied in several rodent species. An increase in the incidence of malignant hepatocellular tumours occurred in male rats and mice following lifetime dietary administration of diazepam at 75 mg/kg/day (17- and 8-fold the MRHD on a body surface area basis, respectively). This was not observed in female rats and mice treated with 75 mg/kg/day or hamsters treated with 120 mg/kg/day (18-fold the MRHD).

Genotoxicity

Limited data from a number of studies have provided weak evidence of a genotoxic potential. Diazepam has been shown to induce aneuploidy in sperm obtained from both mice and humans treated with approximately 10 mg/m²/day (less than the MRHD).

Teratogenicity

Diazepam was found to be teratogenic in mice at intravenous doses of 45 mg/kg or greater and oral doses of 100 mg/kg or greater (both 10-fold the MRHD on a body surface area basis), as well as in hamsters at 280 mg/kg (41-fold the MRHD). The respective no-effect doses were 50 mg/kg (5-fold the MRHD) in mice and 200 mg/kg (30-fold the MRHD) in hamsters. Malformations included exencephaly, cranioschisis, kinking of the spinal cord, and cleft palate with and without cleft lip. Malformations were not observed in rats or rabbits at respective doses of up to 300 and 50 mg/kg/day (greater than 20-fold the MRHD). Delayed development has been reported in offspring from several animal species treated with diazepam during pregnancy or during pregnancy and lactation.

Interactions with Other Medicines

The benzodiazepines, including diazepam, produce additive CNS depressant effects when co-administered with other medications which themselves produce CNS depression e.g. barbiturates, alcohol, anxiolytics, sedatives, antidepressants including tricyclic antidepressants and non-selective MAO inhibitors, hypnotics, antiepileptic drugs, phenothiazines and other antipsychotics, skeletal muscle relaxants, antihistamines, narcotic analgesics and anaesthetics (see PRECAUTIONS). Therefore, it should be borne in mind that the effect of these drugs may potentiate or be potentiated by the action of diazepam.

Concomitant use with alcohol is not recommended due to enhancement of the sedative effect.

There is a potentially relevant interaction between diazepam and compounds that inhibit certain hepatic enzymes, particularly Cytochrome P450 3A (CYP3A). Data indicate that these compounds influence the pharmacokinetics of diazepam and may lead to increased and prolonged sedation. Diazepam undergoes oxidative metabolism and, consequently, may interact with disulfiram, cimetidine, ketoconazole, fluvoxamine, fluoxetine or omeprazole resulting in increased plasma levels of diazepam. Patients should be observed closely for evidence of enhanced benzodiazepine response during concomitant treatment with either disulfiram or cimetidine; some patients may require a reduction in benzodiazepine dosage.

There have also been reports that the metabolic elimination of phenytoin is affected by diazepam.

Cisapride may lead to a temporary increase in the sedative effects of orally administered benzodiazepines due to faster absorption.

The anticholinergic effects of other drugs including atropine and similar drugs, antihistamines and antidepressants may be potentiated.

Interactions have been reported between some benzodiazepines and anticonvulsants, with changes in the serum concentration of the benzodiazepine or anticonvulsant. It is recommended that patients be observed for altered responses when benzodiazepines and anticonvulsants are prescribed together and that serum level monitoring of the anticonvulsant is performed more frequently.

Effects on Laboratory tests

Minor EEG changes, usually low voltage fast activity, of no known clinical significance have been reported with benzodiazepine administration.

Diazepam can inhibit binding of thyroxine and liothyronine to their binding proteins resulting in erroneously abnormal values from thyroid function test.

ADVERSE EFFECTS

Most commonly reported undesirable effects are fatigue, drowsiness, muscle weakness and ataxia; they are usually dose-related.

Effects encountered infrequently are amnesia, confusion, constipation, depression, diplopia, dysarthria, gastrointestinal disturbances, headache, hypotension, incontinence, increase or decrease in libido, nausea, dry mouth or hypersalivation, skin reactions such as rash, slurred speech, tremor, urinary retention, vertigo and blurred vision.

Very rarely, elevated transaminases and alkaline phosphatase, jaundice as well as cases of cardiac arrest have been observed. Isolated instances of neutropenia have been seen. Periodic blood counts and liver function tests are advisable during long-term therapy.

Dizziness has been reported occasionally with diazepam.

Anterograde amnesia may occur using therapeutic dosages, the risk increasing at higher doses. Amnesic effects may be associated with inappropriate behaviour.

Paradoxical reactions such as acute hyperexcitation, anxiety, hallucinations, increased muscle spasticity, insomnia, rage, sleep disturbances and stimulation have been reported; should these occur, use of the drug should be discontinued.

Chronic use (even at therapeutic doses) of oral diazepam may lead to the development of physical dependence: discontinuation of the therapy may result in withdrawal or rebound phenomena.

DOSAGE AND ADMINISTRATION

For maximal beneficial effect, the dosage should be carefully individualised. Dosage may need to be reduced in patients with hepatic or renal disease as the elimination half-life may be prolonged in this subgroup.

Elderly patients should be given a reduced dose. These patients should be checked regularly at the start of treatment in order to minimise the dosage and/or frequency of administration to prevent overdose due to accumulation.

Usual Adult Dosage: 5-40 mg daily.

Average Dosage for Ambulatory Patients: 2 mg three times daily or 5 mg in the evening and 2 mg once or twice during the day.

Elderly or Debilitated Patients: 2 mg twice daily or half the usual adult dose.

Children – 6 months to 3 years: 1-6 mg daily,

Children – 4 to 14 years: 4-12 mg daily or calculated from 0.1-0.3 mg/kg bodyweight.

Hospital treatment of tension, excitation, motor unrest: 10 to 15 mg three times daily until the acute symptoms subside.

Muscle spasm: 10-30 mg daily.

Benzodiazepines should not be given to children without careful assessment of the indication; the duration of treatment must be kept to a minimum.

OVERDOSAGE

Symptoms

Overdosage of benzodiazepines is usually manifested by degrees of central nervous system depression ranging from drowsiness to coma. In mild cases, symptoms include drowsiness, mental confusion and lethargy. In more serious cases, symptoms may include ataxia, hypotonia, hypotension, respiratory depression and rarely, coma and death. Coma, may be more protracted and cyclical, particularly in elderly patients. Benzodiazepine respiratory depressant effects are more serious in patients with respiratory disease.

Benzodiazepines increase the effects of other central nervous system depressants, including alcohol. When combined with other CNS depressants, the effects of overdosage are likely to be severe and may prove fatal.

Treatment

Treatment of overdose is symptomatic; institute supportive measures as indicated by the patient's clinical state. If the overdosage is known to be small, observation of the patient and monitoring of their vital signs only may be appropriate. In adults or children who may have taken an overdose of benzodiazepines within 1-2 hours, consider activated charcoal with airway protection if indicated.

If CNS depression is severe, consider the use of flumazenil (Anexate®), a benzodiazepine antagonist. This should only be administered under closely monitored conditions. It has a short half-life (about an hour), therefore patients administered with flumazenil will require monitoring after its effects have worn off. Flumazenil may precipitate seizures and is contraindicated in the presence of drugs that reduce seizures threshold (e.g. tricyclic antidepressants) and epileptic patients who have been treated with benzodiazepines. Refer to prescribing information for flumazenil (Anexate®), for further information on the correct use of this drug.

Haemoperfusion and haemodialysis are not useful in benzodiazepine intoxication.

Contact the Poisons Information Centre on 13 11 26 (Australia) for advice on the management of overdose.

PRESENTATIONS AND STORAGE CONDITIONS

APO-Diazepam 2 mg Tablets

White to off-white round biconvex uncoated tablets debossed with 2 on one side and plain on the other side.

AUST R number 134472.

APO-Diazepam 5 mg Tablets

White to off-white round flat bevelled edged uncoated tablets debossed with 5 and a scoreline on one side and plain on the other side.

AUST R number 134590

Blister packs of 50 tablets.

APO-Diazepam Tablets are intended for oral administration. Each tablet contains diazepam 2 mg or 5 mg.

In addition, each tablet contains the following inactive ingredients: lactose, maize starch, colloidal anhydrous silica, magnesium stearate and purified talc.

Store below 25°C. Protect from light.

NAME AND ADDRESS OF THE SPONSOR

Apotex Pty Ltd
66 Waterloo Road
North Ryde NSW 2113
Australia

Apotex Pty Ltd is the licensee of the registered trade marks APO and APOTEX from the registered proprietor, Apotex Inc.

POISONS SCHEDULE OF THE MEDICINE:

S4: Prescription Only Medicine

Date of TGA approval: 24 October 2007

Date of most recent amendment: 22 January 2010